

# Snail sni' ta Julbil li poxil Insulina li k'uxi Anil ta Ak'bele (100 yepal/mL)


## Rapid-Acting Insulin Injection Pen (100 units/mL) (Tzotzil)

Sbi li poxile: \_\_\_\_\_

Prescription name: \_\_\_\_\_

K'uxelan li yich'ojtal li sbenta insulina:


Insulin pen parts:



¿Bu stak' ta julba li ta insulina? Where do I inject the insulin?

- Ts'oto ta julbile li bu ak'bil ta sbeta li ta julbile #1-4 ta sjamleje:  
Rotate injections between injection spots #1-4 in the diagram:
  1. **Ta a Ch'ut** (ja'no' ox 2 sjamlej ta joyob a pis) **Stomach** (except for the 2 inches around the belly button)

2. Ta sba tak chiuk ta xokonan a **vo tak**  
(mu me stak li nopol ta sjol a bakane)  
Top and outside of **thighs** (except in the bony area close to the knees)
3. Ta xokonan a **k' obtak** (ja' lek me oy bocho ta sjulot li ta poxil li insulin)  
Back of the **upper arms** (this might work best if someone else is injecting the insulin)
4. **Ta xokonan a pat o chiuk ta sba a chaktak** Lower back or upper buttocks



- Julo ta insulin te ta jk' ol sbektal li ta yolon snukulile, mauk ta yipal.  
Inject your insulin into the layer of fatty tissue under the skin, not into the muscle.
- Mu xa jul ta jun no' ox yabil. Do not inject into the same spot every time.
- Mu xa jul ta bu oy xch' okal chiuk li bu tuch' aneme. Do not inject into moles or scars.

## ¿K'uxi ta julba li ta insulin? How do I inject the insulin?

### Slikeb 1: Cha kel li insulin Step 1: Check the insulin

- Li snail me oy k' ejel ta bu sikubtasbile, lokeso tal bayuk 1-2 sjalil (horas) k' alal li sk' anto xa tunes. If your pen is in the refrigerator, take it out at least 1-2 hours before use.
- K' elo li ta sbi ti bu snail me ja' ti bu cha tunes li insulin. Check the label to make sure you are using the right insulin pen.
- K' elo sk' ak' alil li bakin xlaj xipal li ta 28 k' ak' al tsibtabil tal ta snaile. Check the 28-day expiration date written on the label.

- Lokesbo li smak snaile chiuk k' elo k' uxelan stalel li insulina li bu kejel talel li insulina. Remove the pen cap and check how the insulin looks in the insulin reservoir.
  - Mu xa tunes li insulina me sokeme, muyuk sak o me oy k' usi tik' ile. Do not use insulin that is cloudy, not clear, or has any specks in it.

### **Chibal 2: Cha tikbe li sni'e Step 2: Attach the needle**

- Poko li smak ta uli'e ta jun spokal chiuk alcohol. Wipe the rubber seal with an alcohol pad.
- Lok'esbo li ts'is ta jot sbae li bu smak sni' oy-e. Remove the plastic seal on the end of the outer needle cap.
- Ts'oto o net'o (ja' li k' uxelan chopukal sni'e) li sni'ta stuk'il li bu oy smak ta uli'e li ta sni' snaile. Twist or push (based on the needle type) the needle straight onto the rubber seal of the pen tip.

### **Yoxibal 3: Cha pas bayuk xchanel me leke (o "net'obatel li bu jamal cha vile")**

#### **Step 3: Perform a safety test (or “air shot”)**

#### **Li k'alal ta jujun julele:** For each injection:

- Lok'eso jun batel 2 yepal (cha ts'ot li k' uxelan yepal ja to me la yak a bil 2 li bu ta k'elel li yapale). Select a dose of 2 units (by turning the dosage knob until it is pointing at 2 in the dose window).
- Lokesbo li smak ta sba sni'e (k' ejo) chiuk li smak sni' ta yute (jipo batel ta k'aep). Take off the outer needle cap (save it) and the inner needle cap (throw it away).
- Li sni'sk'eloj muyel ta toyol, majo jutuk li insulina kucha'al slok'an li sboktak myele. With the needle pointing up, tap the insulin reservoir to move the air bubbles to the top.
- Neto li snetubil julele k'alalto ta spajeb chiuk k' elo me oy chlok li insulina ta sni'e. Press the injection button all the way in and make sure insulin comes out of the needle.

- Me muyuk xlok' li insulina, cha' paso ta 2 belta ja' noox ta moj jun sni'. If insulin does not come out, repeat up to 2 more times with the same needle.
- Li insulina me muyuk xlok' ta 3 belta, jelbo li sni' e cha' paso yanbelta. If insulin does not come out after 3 times, change the needle and try again.
- Me k' alal ya jelbe li sni' e muyuk xlok- o li insulina, o yik' al boch' emtale li snaile. If insulin still does not come out after changing the needle, the pen may be broken.

#### **Xchanibal 4: K'elo k'u yepal Step 4: Select your dose**

- K' elo li bu ta xak ta ilele oyuk ta "0" k' alal me la pas me lek abile (jech sbi xtok "cha net' ta jamal"). Check that the dose window shows "0" again after the safety test (also called an "air shot").
- Ts' oto li bu chal sbenta li ku yepal ta xa tunes li k' uselan albiltal yepal li tsibtable. Turn the dosage knob to the number of units that equals your prescribed dose.
  - K' elo ta k' un li mu xa-ta ta netel li bu sbenta julel k' alal li cha kel ku yepale. Be careful not to push the injection button while selecting your dose.

#### **Svobal 5: Cha Jul li k'u yepale Step 5: Inject the dose**

- Kuso li bu yabil cha jul ta jun gasa chiuk alcohol malo k' alal me takije. Clean the injection site with an alcohol swab and wait for it to dry.
- Tik' o ochel li sni' k' alalto ta spajeb li bu snukulile. Put the needle all the way into the skin.
- Net' o chiuk te malo k' u sjalil li snet' ubil ta julel sbenta cha tik' li k' u yepal cha vich' e. Press and hold the injection button to give the dose.
- Te neto' o li snetubil chiuk cha bati ta kun ja' to me la k' ot ta 10 k' alal sk' anto xa lokes li sni' ta a bektale. Keep the button pressed and slowly count to 10 before taking the needle out of the skin.
  - Mu xa pik li bu yabil la jul li ta insulina. Do not rub the spot where you injected the insulin.

## **Sbakibal 6: Lok'eso chiuk cha jip batel li sni'e** Step 6: Remove and discard the needle

- Tik' o sutel li smak ta sba sni'e tuneso sbenta xu' xa lok'es tale li sni' ta snaile. Put the outer needle cap back on the needle and use it to remove the needle from the pen.
  - Cha lokesbe li sni' ta snail k'alal ta ju koj cha tunese. Remove the needle from the pen after each dose.
  - Mu buchouk xa bak' be ta yantik li sni' chiuk li snail li insulina. Do not share needles or insulin pens with anyone else.
  - Mu xa cha tunes li sni'take. Do not re-use needles.
- Tik' o ochel ta jun tsoj snail li sni' bu a tunesoje ja li bu akbilot sbenta cha kejan li bu oyay snitake, me oy bu yan snail xu' xa kej li oyuk smoke (jech kucha' al jun yabil shabon bu xokol la tunes ta a k'utake). Place the used needle in the red sharps container we gave you, or in a hard plastic container with a sealable lid (like an empty laundry detergent bottle).
  - Kejbo li snail li insulina muyuk tik' il li sni'e chiuk makaluk lek. Store the insulin pen without a needle attached and with the pen cap on.

## **¿Bu xu' jk'ej li snail li insulina? Where do I store the insulin pen?**

- Xu' xa k'ejanan li snailtak li insulina li bu muyuk pikbil ta jun sikubtasbil (refrigerador). You can keep unused insulin pens in the refrigerator.
- Li snailtak li bu la pikaxae xu' xa bak ta bu lek no' ox k'ixnale ta sjalil 28 k'ak' ale k'alal ti la lik a tunesane. Keep in-use pens at room temperature for no more than 28 days once you begin using them.
- Me jelo li 28 k'ak' ale, jipo xa batel ta k'aep li sailan insulina li bu a tunesanojxae, me oy to jutuk li insulina te xa jip-o batel. After 28 days, throw away in-use insulin pens in the trash, even if there is leftover insulin in the pen.

- Mu xa k' ejan li snail insulina li bu k' ixin yabiltak o me ta xta' at jutuk ta xojabal jtotik. Do not keep insulin pens where there is direct heat or sunlight on the pen.
- Li snailtak li insulina oy patsubenan ta sikile, jipoxa batelan. If the insulin pens are frozen or overheated, throw them away.

### **Liol tsotsik sbenta k'uxelan ta tunesel li snail insulina: Important tips for using the insulin pen:**

- Tuneso li insulina sbenta li ta anillta akbele kala me muyem up li yasukar ta chichale chiuk li carbihidratos ja kuxelan albiloxuk tuun li jpoxtabaneje. Use the rapid-acting insulin for high blood sugars and carbohydrates as directed by your medical team.
- Jelano li yabiltak li bu 4 albil sbenta stak julelanel. Rotate your injections between the 4 injection spots listed above.
- Ta juju ten tuneso ach li sni' e chiuk kuso lek li bu yabil cha julane. Always use a new needle for each injection, and clean the injection site before each dose.
- Atio ta k' un k' alalto ta 10 k' alal me la net li ku jepale cha tike k' alal li sk' anto xa lokes li sni' ta a nukulile. Slowly count to 10 after pressing the injection button before you remove the needle from the skin.
- Me oy lok li insulina ta a nukuli k' alal la jule, mu xa bak yan. Ja no' ox na' olek yatiel li 10 k' alal tik' il li sni' ta a nukil ta yan jkojele. If insulin leaks out of your skin after the injection, do not give another dose. Just be sure to keep the needle in the skin for at least 10 seconds next time.

Ta slok ta jba: Ti li' bune yichoxtalel loil ayej chiuk/o k'usi talem smelolal pasbil yu'un  
University of Michigan (U-M) Health ta sbenta li bucho ip chiuk ta sbenta skelel. Oy  
yichanoj tale li bu stak túnel ta anil ta línea li bu muyuk pasanbil yun U-M Health chiukan li  
U-M Health muyuk chak ta sba. Mu xa jel li chanubtasbanej ta poxtabaneje li bocho oy yan  
ta skelote, ya ti mu mojuk xchanojanik li k'uxelan ta pasanel ta k'elel li bocho ipike.  
Chi'no ta loil li bocho ta skoltaot li ta poxtabaneje me oy k'usi cha k'an cha jak'an  
sbenta li bune, sbenta ta k'elel chiuk li skolele.

Disclaimer: This document contains information and/or instructional materials developed by University of Michigan (U-M) Health for the typical patient with your condition. It may include links to online content that was not created by U-M Health and for which U-M Health does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition, or your treatment plan.

Sbitak: Jen Kelley, MSN RN WTA-C CPN ACCNS-P, Laura Hailes, BSN RN CPN, Kelly Ruemenapp, BSN RN CPN

Authors: Jen Kelley, MSN RN WTA-C CPN ACCNS-P, Laura Hailes, BSN RN CPN, Kelly Ruemenapp, BSN RN CPN

Cha meltsanbil: Brittany Batell, MPH MSW  
Edited by: Brittany Batell, MPH MSW

Li bocho' yakojtal lokombail: "[Parts of an Insulin Pen](#)" ta Hesham Elsayed (li bu la xich' taele "[Evaluation of Subcutaneous Glucose Monitoring Systems under Routine Environmental Conditions](#)" sveal ResearchGate)

Image attribution: "[Parts of an Insulin Pen](#)" by Hesham Elsayed (source publication "[Evaluation of Subcutaneous Glucose Monitoring Systems under Routine Environmental Conditions](#)" via ResearchGate)

Patient Education ta [U-M Health](#) oy yichoxtalel sk'elel ta [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#). Slajem lok' em 03/2024

Patient Education by [U-M Health](#) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International Public License](#). Last revised 03/2024