

How to Use Your Breyna™ Inhaler (budesonide/formoterol) for Asthma Anti-Inflammatory Reliever (AIR) Therapy

Breyna[™] inhaler (80/4.5 mcg or 160/4.5 mcg)



What is the purpose of this medication?

Breyna[™] is an inhaled corticosteroid and long-acting beta-2 agonist that helps you breathe easier and improves your lung function.

What is asthma anti-inflammatory reliever (AIR) therapy?

Asthma AIR therapy is a new strategy to better manage asthma when patients have an asthma flare-up (also called an asthma attack). It involves an increase of inhaled corticosteroid medication along with a beta-agonist medication to treat inflammation (swelling) and bronchospasms (tightening of the airways in your lungs). In some cases, your regular asthma medication may be the same medication used for AIR therapy because it works quickly.

How and when should I prime this medication?

- If your Breyna[™] inhaler is new, or if you have not used it in 7 days, you will need to **prime** your inhaler. Priming an inhaler means making sure the medication is mixed with the spray so you will get the correct dose of medication when you use the inhaler.
- Prime your inhaler by removing the mouthpiece cover, shaking the inhaler for 5 seconds, and spraying it once away from you. Repeat these steps for a total of 4 sprays.

How do I use my Breyna[™] inhaler for asthma AIR therapy?

- 1. Remove the mouthpiece cover. Put your middle or index finger on top of the canister and put your thumb under the mouthpiece of the inhaler.
- 2. Shake your inhaler for 5-10 seconds.
 - If you're able to use a spacer, or if your doctor has recommended you use a spacer, attach the spacer to your inhaler after shaking it and then remove the mouthpiece cap from your spacer.
- 3. Breathe out fully (to prepare to breathe in the medication).
- 4. Put the mouthpiece in your mouth and close your lips around it. Do not block the mouthpiece with your teeth or tongue.
 - If you're using a spacer, make sure to hold the spacer with the inhaler upright and tilt your head back slightly when you close your lips around the spacer's mouthpiece.
- 5. Push the top all the way down one time while breathing in deeply and slowly through your mouth.
 - If you're using a spacer, push the canister down to release the medication into the spacer. Then breathe in slowly and deeply through the spacer mouthpiece.
- 6. Hold your breath for 5-10 seconds, and then breathe out.
 - If your doctor told you to take more than 1 dose (puff), wait 30

Pulmonary and Critical Care Medicine

How to Use Your Breyna™ Inhaler (budesonide/formoterol) for Asthma Anti-Inflammatory Reliever (AIR) Therapy seconds and then repeat the steps above.

 Rinse your mouth well with water after taking this medication to decrease the risk of a mouth infection. Spit out the water you rinsed with (do not swallow it).

How do I clean my Breyna[™] inhaler?

Clean your inhaler once per week to prevent the medication from building up and blocking the spray.

- 1. Remove the mouthpiece cover.
- 2. Use a clean, dry cotton swab (Q-tip®) to clean the small circular opening where the medication sprays out of the canister.
- 3. Wipe the inside of the mouthpiece with a little bit on water on a clean tissue.
- 4. Let your inhaler air dry.
- 5. Put the cover back over the mouthpiece after it has dried.

How should I store my Breyna[™] inhaler?

Store your inhaler in a cool, dry place away from moisture, heat, or sunlight.

Where can I learn more about my condition or inhaler?

- To learn more about your health condition, visit: <u>CareGuides.med.umich.edu</u>
- To learn more about the Breyna[™] inhaler, visit: <u>Breyna.com</u>

Disclaimer: This document contains information and/or instructional materials developed by University of Michigan (U-M) Health for the typical patient with your condition. It may include links to online content that was not created by U-M Health and for which U-M Health does not assume responsibility. It does not replace medical advice from your health care provider because your experience may differ from that of the typical patient. Talk to your health care provider if you have any questions about this document, your condition, or your treatment plan.

Author: Arjun Mohan, MD Edited by: Brittany Batell, MPH MSW

Patient Education by <u>U-M Health</u> is licensed under a <u>Creative Commons Attribution</u> <u>NonCommercial-ShareAlike 4.0 International Public License</u>. Last revised 05/2024