

Supplement to Nair et al..

Localization of Psoriasis Susceptibility Locus PSORS1 to a 60 Kilobase Interval Telomeric to HLA-C.

HLA-Cw6 phenotype frequencies for psoriatic individuals in world populations.

Population	Frequency (%)	Relative Risk	Reference
Brazilians	59	10	Gonzaga et al., 1996
British	47	3.5	Mallon et al., 1997
Chinese	17	19.8	Cao et al., 1993
Danes	47	12.9	Marcusson et al., 1981
Finns	46	11.5	Tiillikainen et al., 1980
Germans	57	5.3	Hohler et al., 1996
Greeks	36	3.3	Economidou et al., 1985
Israelis	25	1.5	Roitberg-Tambur et al., 1994
Italians	33	3.5	Nini at al., 1989
Japanese	8-28	12.7-8.2	Nakagawa at al., 1991; Okhido et al., 1982
Spaniards	18	6.2	Gonzalez et al., 1999

References

- Cao K, Song FJ, Li HG, Xu SY, Liu ZH, Su XH, Wang FX (1993) Association between HLA antigens and families with psoriasis vulgaris. Chin Med J 106:132-5
- Economidou J, Papasteriades C, Varla-Leftherioti M, Vareltzidis A, Stratigos J (1985) Human lymphocyte antigens A, B, and C in Greek patients with psoriasis: relation to age and clinical expression of the disease. J Am Acad Dermatol 13:578-82
- Gonzaga HF, Torres EA, Alchorne MM, Gerbase Delima M (1996) Both psoriasis and benign migratory glossitis are associated with HLA-Cw6. Br J Dermatol 135:368-70
- Gonzalez S, Martinez-Borra J, Torre-Alonso JC, Gonzalez-Roces S, Sanchez del Rio J, Rodriguez Perez A, Brautbar C, et al (1999) The MICA-A9 triplet repeat polymorphism in the transmembrane region confers additional susceptibility to the development of psoriatic arthritis and is independent of the association of Cw*0602 in psoriasis. Arthritis Rheum 42:1010-6
- Hohler T, Weinmann A, Schneider PM, Rittner C, Schopf RE, Knop J, Hasenclever P, et al (1996) TAP-polymorphisms in juvenile onset psoriasis and psoriatic arthritis. Hum Immunol 51:49-54
- Mallon E, Bunce M, Wojnarowska F, Welsh K (1997) HLA-CW*0602 is a susceptibility factor in type I psoriasis, and evidence Ala-73 is increased in male type I psoriatics. J Invest Dermatol 109:183-6
- Marcusson JA, Johannesson A, Moller E (1981) HLA-A,B,C and DR antigens in psoriasis. Tissue Antigens 17:525-9
- Nakagawa H, Akazaki S, Asahina A, Tokunaga K, Matsuki K, Kuwata S, Ishibashi Y, et al (1991) Study of HLA class I, class II and complement genes (C2, C4A, C4B and BF) in Japanese psoriatics and analysis of a newly-found high-risk haplotype by pulsed field gel electrophoresis. Arch Dermatol Res 283:281-4
- Nini G, Bianchi L, Iraci S, Camplone G, Spagnuolo A, Adorno D, Papola F (1989) HLA antigens and infantile psoriasis. Acta Derm Venereol Suppl 146:59-62
- Okhido M, Ozawa A, Nakagawa M, Tsuji K (1982) HLA-Cw6 in Japanese psoriasis vulgaris. In: Farber EM, Cox AJ (eds) Psoriasis: Proceedings of the Third International Symposium. Grune & Stratton, New York, pp 287-288
- Roitberg-Tambur A, Friedmann A, Tzfoni EE, Battat S, Ben Hammo R, Safirman C, Tokunaga K, et al (1994) Do specific pockets of HLA-C molecules predispose Jewish patients to psoriasis vulgaris? J Am Acad Dermatol 31:964-8

Tiilikainen A, Lassus A, Karvonen J, Vartiainen P, Julin M (1980) Psoriasis and HLA-Cw6. Br J Dermatol 102:179-84